

ON THE TRACE OF HECKE OPERATORS FOR MAASS FORMS FOR CONGRUENCE SUBGROUPS II

XIAN-JIN LI

ABSTRACT. Let E_λ be the Hilbert space spanned by the eigenfunctions of the non-Euclidean Laplacian associated with a positive discrete eigenvalue λ . In this paper, the trace of Hecke operators T_n acting on the space E_λ is computed for Hecke congruence subgroups $\Gamma_0(N)$ of non-square free level. This extends the computation of Conrey-Li [2], where only Hecke congruence subgroups $\Gamma_0(N)$ of square free level N were considered.

1. INTRODUCTION

Let $N > 1$ be an integer, which is not square free. Denote by $\Gamma_0(N)$ the Hecke congruence subgroup of level N . The non-Euclidean Laplacian Δ on the upper half-plane $\mathcal{H}$ is given by

$$\Delta = -y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right).$$

Let D be the fundamental domain of $\Gamma_0(N)$. Eigenfunctions of the discrete spectrum of Δ are nonzero real-analytic solutions of the equation

$$\Delta\psi = \lambda\psi$$

such that $\psi(\gamma z) = \psi(z)$ for all γ in $\Gamma_0(N)$ and such that

$$\int_D |\psi(z)|^2 dz < \infty$$

where dz represents the Poincaré measure of the upper half-plane.

Let $\mathfrak{a}$ be a cusp of $\Gamma_0(N)$. We denote its stabilizer by $\Gamma_{\mathfrak{a}}$. An element $\sigma_{\mathfrak{a}}$ in $PSL_2(\mathbb{R})$ exists such that $\sigma_{\mathfrak{a}}\infty = \mathfrak{a}$ and $\sigma_{\mathfrak{a}}^{-1}\Gamma_{\mathfrak{a}}\sigma_{\mathfrak{a}} = \Gamma_\infty$. If ψ is an eigenfunction of the Laplacian associated with a positive discrete eigenvalue λ , then it has a Fourier expansion [6]

$$\psi(\sigma_{\mathfrak{a}}z) = \sqrt{y} \sum_{m \neq 0} \rho_{\mathfrak{a}}(m) K_{i\kappa}(2\pi|m|y) e^{2m\pi ix}$$

1991 *Mathematics Subject Classification.* Primary 11F37, 11F72, 42A16.

Key words and phrases. Hecke operators, Laplace-Beltrami operator, Maass wave forms, Selberg trace formula.

Research partially supported by the American Institute of Mathematics.

at every cusp $\mathfrak{a}$ of $\Gamma_0(N)$, where $\kappa = \sqrt{\lambda - 1/4}$ and $K_\nu(y)$ is given by the formula §6.32, [18]

$$(1.1) \quad K_\nu(y) = \frac{2^\nu \Gamma(\nu + \frac{1}{2})}{y^\nu \sqrt{\pi}} \int_0^\infty \frac{\cos(yt)}{(1+t^2)^{\nu+\frac{1}{2}}} dt.$$

The Hecke operators T_n , $n = 1, 2, \dots$, $(n, N) = 1$, which act in the space of automorphic functions with respect to $\Gamma_0(N)$, are defined by

$$(T_n f)(z) = \frac{1}{\sqrt{n}} \sum_{ad=n, 0 \leq b < d} f\left(\frac{az+b}{d}\right).$$

An orthonormal system of eigenfunctions of Δ exists [6] such that each of them is an eigenfunction of all the Hecke operators. We call these eigenfunctions Maass-Hecke eigenfunctions. Let λ_j , $j = 1, 2, \dots$, be an enumeration in increasing order of all positive discrete eigenvalues of Δ for $\Gamma_0(N)$ with an eigenvalue of multiplicity m appearing m times, and let $\kappa_j = \sqrt{\lambda_j - 1/4}$. If $\psi_j(z)$ is a Maass-Hecke eigenfunction of Δ associated with the j th eigenvalue λ_j , then

$$(T_n \psi_j)(z) = \tau_j(n) \psi_j(z)$$

where $\rho_{j\infty}(m) = \rho_{j\infty}(d) \tau_j(n)$ if $m = dn$ with $n \geq 1$, $(n, dN) = 1$.

Let E_λ be a Hilbert space of functions spanned by the eigenfunctions of Δ associated with a positive discrete eigenvalue λ . The Petersson inner product of the space is given by

$$(1.2) \quad \langle F(z), G(z) \rangle = \int_D F(z) \bar{G}(z) dz.$$

The analogue for Maass forms of Eichler-Selberg's trace formula [12], p.85 is obtained for congruence subgroups $\Gamma_0(N)$ of square free level N in Conrey-Li [2]. In this paper, the trace $\text{tr}_\lambda T_n$ of Hecke operators acting on the space E_λ is computed for congruence subgroups $\Gamma_0(N)$ of non-square free N . Some of this computation is implicit in Hejhal [4][5].

Denote by h_d the class number of indefinite rational quadratic forms with discriminant d . Let

$$(1.3) \quad \epsilon_d = \frac{v_0 + u_0 \sqrt{d}}{2},$$

where the pair (v_0, u_0) is the fundamental solution [10] of Pell's equation $v^2 - du^2 = 4$. Denote by Ω the set of all the positive integers d such that $d \equiv 0$ or $1 \pmod{4}$ and such that d is not a square of an integer. Let $\mu(n)$ be the Möbius function.

Theorem 1. *Let N be a non-square free positive integer, and let n be a positive integer with $(n, N) = 1$. If*

$$L_n(s) = \sum_{k|N} k^{1-2s} \sum_{m|\frac{N}{k}} \sum_{n|k} \frac{\mu(n)}{n} \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \left(\frac{d}{mn} \right) \frac{h_d \ln \epsilon_d}{(du^2)^s}$$

for $\text{Re } s > 1$, where the sum on u is over all the positive integers u such that $\sqrt{4n + dk^2u^2} \in \mathbb{Z}$ and the product on p^{2l} runs over all distinct primes p with p^{2l} being the greatest even p -power factor of $(d, \frac{N}{k})$, then $L_n(s)$ is analytic for $\text{Re } s > 1$ and is extended by analytic continuation to the half-plane $\text{Re } s > 0$ except for a possible pole at $s = 1/2$ and for possible simple poles at $s = 1, \frac{1}{2} \pm i\kappa_j$, $j = 1, 2, \dots$. Moreover, we have

$$\text{tr}_{\lambda_j} T_n = 2n^{i\kappa_j} \text{Res}_{s=1/2+i\kappa_j} L_n(s)$$

for $j = 1, 2, \dots$.

The paper is organized as follows. In section 2, we recall the Selberg trace formula; see Hejhal [4] [5] and Selberg [12]. Then in section 3 we compute contributions of identity, elliptic, hyperbolic and parabolic elements to the trace formula. A technical part of this section is to compute the Petersson inner product of an Eisenstein series for $\Gamma_0(N)$ with the image of the Eisenstein series under the action of Hecke operators. The result is given in Theorem 3.11. By using the Selberg trace formula, we obtain analyticity information about a series formed by contributions to the trace formula of hyperbolic elements whose fixed points are not cusps. A precise statement is given in Theorem 3.13. In section 4, we compute explicitly the total contribution to the trace formula of hyperbolic elements whose fixed points are not cusps. The result is stated in Theorem 4.6. Finally, Theorem 1 follows from Theorem 3.13 and Theorem 4.6.

Most part of this paper was written at the American Institute of Mathematics, California. The author wishes to thank AIM and Brian Conrey for the support. He also wishes to thank Dennis Hejhal and Henryk Iwaniec for helpful conversations and for their encouragement for the proof of Theorem 3.11.

2. THE SELBERG TRACE FORMULA

Assume that s is a complex number with $\text{Re } s > 1$. Let

$$k(t) = \left(1 + \frac{t}{4}\right)^{-s}$$

and

$$k(z, z') = k\left(\frac{|z - z'|^2}{yy'}\right),$$

for $z = x + iy$ and $z' = x' + iy'$ in the upper half-plane. The kernel function $k(z, z')$ is of (a)-(b) type in the sense of Selberg [12], p.60. Let

$$g(u) = \int_w^\infty k(t) \frac{dt}{\sqrt{t-w}}$$

with $w = e^u + e^{-u} - 2$. If

$$h(r) = \int_{-\infty}^{\infty} g(u) e^{iru} du,$$

then

$$(2.1) \quad g(u) = c(1 + \frac{w}{4})^{\frac{1}{2}-s}$$

where $c = 2\sqrt{\pi}\Gamma(s - \frac{1}{2})\Gamma^{-1}(s)$. Since

$$h(r) = c \frac{4^s(s - \frac{1}{2})}{(s - \frac{1}{2})^2 + r^2} + A(r, s),$$

where $A(r, s)$ is finite for $|\Im r| \leq 1/2$ and for $\operatorname{Re} s > 0$, we find that

$$(2.2) \quad \lim_{s \rightarrow 1/2 + i\kappa} (s - \frac{1}{2} - i\kappa)h(r) = \begin{cases} 4^{1/2+i\kappa} \sqrt{\pi} \frac{\Gamma(i\kappa)}{\Gamma(1/2+i\kappa)}, & \text{for } r = \pm\kappa; \\ 0, & \text{for } r \neq \pm\kappa. \end{cases}$$

Assume that n is a positive integer with $(n, N) = 1$. Let

$$\Gamma^* = \cup_{\substack{ad=n \\ 0 \leq b < d}} \frac{1}{\sqrt{n}} \begin{pmatrix} d & -b \\ 0 & a \end{pmatrix} \Gamma_0(N).$$

Then Γ^* satisfies all the requirements given in [12], p.69. Let $\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_{\nu(N)}$ be a complete set of inequivalent cusps of $\Gamma_0(N)$. We choose $\sigma_{\mathfrak{a}_i} \in PSL_2(\mathbb{R})$ such that $\sigma_{\mathfrak{a}_i} \infty = \mathfrak{a}_i$ and $\sigma_{\mathfrak{a}_i}^{-1} \Gamma_{\mathfrak{a}_i} \sigma_{\mathfrak{a}_i} = \Gamma_{\infty}$ for $i = 1, 2, \dots, \nu(N)$. Let $E_i(z, s)$ be the Eisenstein series for the cusp $\mathfrak{a}_i$, let

$$K(z, z') = \sum_{T \in \Gamma^*} k(z, Tz'),$$

and let

$$H(z, z') = \sum_{i=1}^{\nu(N)} \sum_{ad=n, 0 \leq b < d} \frac{1}{4\pi} \int_{-\infty}^{\infty} h(r) E_i(\frac{az+b}{d}, \frac{1}{2} + ir) E_i(z', \frac{1}{2} - ir) dr.$$

It follows from (2.14) of [12], the argument of [7], pp.96-98, Theorem 5.3.3 of [7], and the spectral decomposition formula (5.3.12) of [7] that

$$(2.3) \quad d(n)h(-\frac{i}{2}) + \sqrt{n} \sum_{j=1}^{\infty} h(\kappa_j) \operatorname{tr}_{\lambda_j} T_n = \int_D \{K(z, z) - H(z, z)\} dz$$

for $\operatorname{Re} s > 1$, where $d(n)$ is the sum of positive divisors of n .

3. EVALUATION OF COMPONENTS OF THE TRACE

For every element T of Γ^* , we denote by Γ_T the set of all the elements of $\Gamma_0(N)$ which commute with T . Let $D_T = \Gamma_T \backslash \mathcal{H}$. Elements of Γ^* can be classified as the identity, the hyperbolic, the elliptic, and the parabolic elements. If T is not a parabolic element, we put

$$c(T) = \int_{D_T} k(z, Tz) dz.$$

Proofs for the following lemmas from Lemma 3.1 to Lemma 3.8 are minor modifications of corresponding lemmas in Conrey-Li [2]. For the convenience of readers, we sketch the proofs here.

3.1. The identity component.

If Γ^* contains the identity element I , then

$$c(I) = \int_{\Gamma_0(N) \backslash \mathcal{H}} dz.$$

3.2. Elliptic components.

There are only a finite number of elliptic conjugacy classes.

Lemma 3.1. *Let R be an elliptic element of Γ^* . Then*

$$c(R) = \frac{\pi}{2m \sin \theta} \int_0^\infty \frac{k(t)}{\sqrt{t + 4 \sin^2 \theta}} dt,$$

where m is the order of a primitive element of Γ_R and where θ is defined by the formula $\text{trace}(R) = 2 \cos \theta$.

Proof. Since R is an elliptic element of Γ^* , an element $\sigma \in PSL(2, \mathbb{R})$ exists such that

$$\sigma R \sigma^{-1} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} = \tilde{R}$$

for some number $0 < \theta < \pi$. If $(\sigma \Gamma_0(N) \sigma^{-1})_{\tilde{R}}$ is the set of all the elements of $\sigma \Gamma_0(N) \sigma^{-1}$ commuting with $\tilde{R}$, then we have

$$c(R) = \int_{D_{\tilde{R}}} k(z, \tilde{R}z) dz$$

where $D_{\tilde{R}} = (\sigma \Gamma_0(N) \sigma^{-1})_{\tilde{R}} \backslash \mathcal{H}$.

If γ is an element of $\Gamma_0(N)$ having the same fixed points as R , then γ commutes with R . By Proposition 1.16 of [13], a primitive elliptic element γ_0 of $\Gamma_0(N)$ exists such that $(\eta \Gamma \eta^{-1})_{\tilde{R}}$ is generated by $\eta \gamma_0 \eta^{-1}$. Since $\eta \gamma_0 \eta^{-1}$ commutes with $\tilde{R}$, it is of the form

$$\begin{pmatrix} \cos \theta_0 & -\sin \theta_0 \\ \sin \theta_0 & \cos \theta_0 \end{pmatrix}$$

for some real number θ_0 . By Proposition 1.16 of [13], $\theta_0 = \pi/m$ for some positive integer m . It follows from the argument of [7], p.99 that

$$c(R) = \frac{1}{m} \int_0^\infty \int_{-\infty}^\infty k \left(\frac{|z^2 + 1|^2}{y^2} \sin^2 \theta \right) dz.$$

By the argument of [7], p.100 we have

$$c(R) = \frac{\pi}{2m \sin \theta} \int_0^\infty \frac{k(t)}{\sqrt{t + 4 \sin^2 \theta}} dt. \quad \square$$

3.3. Hyperbolic components.

Let P be a hyperbolic element of Γ^* . Then an element ρ exists in $SL_2(\mathbb{R})$ such that

$$\rho P \rho^{-1} = \begin{pmatrix} \lambda_P & 0 \\ 0 & \lambda_P^{-1} \end{pmatrix} = \tilde{P}$$

with $\lambda_P > 1$. The number λ_P^2 is called the norm of P , and is denoted by NP . It follows that

$$c(P) = \int_{D_{\tilde{P}}} k(z, NPz) dz$$

where $D_{\tilde{P}} = (\rho \Gamma_0(N) \rho^{-1})_{\tilde{P}} \backslash \mathcal{H}$. Let P_0 be a primitive hyperbolic element of $SL_2(\mathbb{Z})$, which generates the group of all elements of $SL_2(\mathbb{Z})$ commuting with P . Then there exists a “primitive” hyperbolic element $P_1 \in \Gamma_0(N)$, which generates Γ_P , such that P_1 is the smallest positive integer power of P_0 among all the generators of Γ_P in $\Gamma_0(N)$.

Lemma 3.2. *Let P be a hyperbolic element of Γ^* such that $\Gamma_P \neq \{1_2\}$. If P_1 is a “primitive” hyperbolic element of $\Gamma_0(N)$ which generates the group Γ_P , then*

$$c(P) = \frac{\ln NP_1}{(NP)^{1/2} - (NP)^{-1/2}} g(\ln NP).$$

Proof. Since $\rho P_1 \rho^{-1}$ commutes with $\tilde{P}$, it is of the form

$$\begin{pmatrix} \lambda_{P_1} & 0 \\ 0 & \lambda_{P_1}^{-1} \end{pmatrix}$$

for some number $\lambda_{P_1} > 1$, and hence

$$c(P) = \int_1^{NP_1} \frac{dy}{y^2} \int_{-\infty}^\infty k \left(\frac{(NP - 1)^2 |z|^2}{NP y^2} \right) dx.$$

The stated identity then follows. $\square$

Lemma 3.3. *Let $P = \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ be a hyperbolic element of Γ^* such that $\Gamma_P = \{1_2\}$. Then fixed points of P are cusps of $\Gamma_0(N)$. Moreover, $\Gamma_P = \{1_2\}$ if and only if $A + D = \frac{1}{2}(m + \frac{4n}{m})$ for some divisor m of $4n$ with $m \neq 2\sqrt{n}$ for $C \neq 0$, and if and only if $A \neq D$ for $C = 0$.*

Proof. If the fixed points of P are not rational numbers or infinity, then they are zeros of an irreducible polynomial $ax^2 + bx + c$ with $a \equiv 0 \pmod{N}$. If $d = b^2 - 4ac$ and (u, v) is a solution of Pell's equation $v^2 - du^2 = 4$, then

$$\begin{pmatrix} \frac{v-bu}{2} & -cu \\ au & \frac{v+bu}{2} \end{pmatrix}$$

belongs to $\Gamma_0(N)$ and commutes with P . This contradicts to $\Gamma_P = \{1_2\}$. Hence, fixed points of P are cusps of $\Gamma_0(N)$.

Conversely, if T is an element of Γ^* having two distinct fixed points with at least one of them being a rational number, then we must have $\Gamma_T = \{1_2\}$. Hence, $\Gamma_P = \{1_2\}$ for an element $P = \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma^*$ if and only if the fixed points of P are rationals or the infinity, that is, if and only if $A + D = \frac{1}{2}(m + \frac{4n}{m})$ for some divisor m of $4n$ with $m \neq 2\sqrt{n}$ if $C \neq 0$ or $A \neq D$ if $C = 0$. $\square$

Every cusp of $\Gamma_0(N)$ is equivalent to one of the following inequivalent cusps

$$(3.1) \quad \frac{u}{w} \text{ with } u, w > 0, (u, w) = 1, w|N.$$

Two such cusps u/w and u_1/w_1 are $\Gamma_0(N)$ -equivalent if and only if $w = w_1$ and $u \equiv u_1 \pmod{(w, N/w)}$. Let $\mathfrak{a} = u/w$ be given as in (3.1). By (2.2) and (2.3) of Deshouillers and Iwaniec [3], we have

$$(3.2) \quad \Gamma_{\mathfrak{a}} = \left\{ \begin{pmatrix} 1 + cu/w & -cu^2/w^2 \\ c & 1 - cu/w \end{pmatrix} : c \equiv 0 \pmod{[w^2, N]} \right\}$$

and

$$\sigma_{\mathfrak{a}}\infty = \mathfrak{a} \text{ and } \sigma_{\mathfrak{a}}^{-1}\Gamma_{\mathfrak{a}}\sigma_{\mathfrak{a}} = \Gamma_{\infty}$$

where

$$(3.3) \quad \sigma_{\mathfrak{a}} = \begin{pmatrix} \mathfrak{a}\sqrt{[w^2, N]} & 0 \\ \sqrt{[w^2, N]} & 1/\mathfrak{a}\sqrt{[w^2, N]} \end{pmatrix}.$$

Let P be a hyperbolic element of Γ^* such that $\Gamma_P = \{1_2\}$. Assume that $\mathfrak{a}$ is a fixed point of P . Then ∞ is a fixed point of $\sigma_{\mathfrak{a}}^{-1}P\sigma_{\mathfrak{a}}$, and hence there exist positive integers a, d with $ad = n, a \neq d$ such that

$$(3.4) \quad P = \frac{1}{\sqrt{n}}\sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1} = \frac{1}{\sqrt{n}} \begin{pmatrix} a - b[w^2, N]\frac{u}{w} & b[w^2, N]\frac{u^2}{w^2} \\ (a - d)\frac{w}{u} - b[w^2, N] & d + b[w^2, N]\frac{u}{w} \end{pmatrix}.$$

Since $c(P)$ depends only on the conjugacy class $\{P\}$ represented by P , we can replace P by $\gamma^{-1}P\gamma$ without changing the value of $c(P)$. Replacing P by $\gamma^{-1}P\gamma$ for some element $\gamma = \sigma_{\mathfrak{a}} \begin{pmatrix} 1 & \ell \\ 0 & 1 \end{pmatrix} \sigma_{\mathfrak{a}}^{-1} \in \Gamma_{\mathfrak{a}}$, we can assume without loss of generality that $0 \leq b < |a - d|$ in (3.4).

Lemma 3.4. *Let $\mathfrak{a} = u/w$ be given as in (3.1). Then*

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

with $a, d \in \mathbb{Z}, ad = n, a \neq d$ is a hyperbolic element of Γ^* with $\Gamma_P = \{1_2\}$ and $P(\mathfrak{a}) = \mathfrak{a}$ if and only if $(w, N/w) | (a - d)$ with b being chosen so that $P \in \Gamma^*$.

Proof. Let P be a hyperbolic element of Γ^* with $\Gamma_P = \{1_2\}$ and $P(\mathfrak{a}) = \mathfrak{a}$. By (3.4), there exists an integer k such that $\frac{u}{w}N | a - d - kw$, and hence $(w, N/w) | a - d$.

Conversely, if a, d are positive integers satisfying $ad = n, a \neq d$ and $(w, N/w) | a - d$, then $a - d = \ell(w, N/w)$ for some integer ℓ . Integers λ and τ exist such that $\lambda \mathfrak{a}N + \tau w = (w, N/w)$, and hence $\ell \lambda \mathfrak{a}N = a - d - \ell \tau w$. Thus, an integer k exists such that $\frac{u}{w}N | a - d - kw$. Let $A = a - kw, B = ku, C = \frac{w}{u}(a - d - kw)$ and $D = d + kw$. Then $P = \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is an element of Γ^* with $\Gamma_P = \{1_2\}$ and $P(\mathfrak{a}) = \mathfrak{a}$, which is of the form

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

with $a, d \in \mathbb{Z}, ad = n, a \neq d$ and with b being chosen so that $P \in \Gamma^*$. $\square$

Lemma 3.5. *Let $\mathfrak{a} = u/w$ be given as in (3.1). Let $a, d \in \mathbb{Z}^+, ad = n, a \neq d$. If $(w, N/w) | a - d$, then there are exactly $|a - d|$ number of $\Gamma_0(N)$ -inequivalent hyperbolic elements $P \in \Gamma^*$ with $P(\mathfrak{a}) = \mathfrak{a}$ and $\Gamma_P = \{1_2\}$, which are of the form*

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

with $0 \leq b < |a - d|$.

Proof. For $a, d \in \mathbb{Z}, ad = n, a \neq d$, let

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1} \quad \text{and} \quad P' = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b' \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

be two elements of Γ^* with $0 \leq b, b' < |a - d|$ whose fixed points are cusps. By using (3.4), we can find that $b' = b + \ell$ for some integer ℓ .

Conversely, for a given integer ℓ with $0 \leq b + \ell < |a - d|$, if $b' = b + \ell$ then

$$P' = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b' \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

is a hyperbolic element of Γ^* with $\Gamma_{P'} = \{1_2\}$ and $P'(\mathfrak{a}) = \mathfrak{a}$. Moreover, if $b' \neq b$ modulo $|a - d|$, then P' and P are not $\Gamma_0(N)$ -equivalent. Therefore, for fixed $a, d \in \mathbb{Z}^+, ad = n, a \neq d$, there are exactly $|a - d|$ $\Gamma_0(N)$ -inequivalent hyperbolic elements P in Γ^* with $P(\mathfrak{a}) = \mathfrak{a}$ and $\Gamma_P = \{1_2\}$, which are of the form

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

with $0 \leq b < |a - d|$. $\square$

Lemma 3.6. *Let $\mathfrak{a} = u/w$, and let*

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1} \quad \text{and} \quad P' = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} d & b' \\ 0 & a \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

be two hyperbolic elements of Γ^ with $P(\mathfrak{a}) = \mathfrak{a}$, $P'(\mathfrak{a}) = \mathfrak{a}$, $\Gamma_P = \{1_2\}$ and $\Gamma_{P'} = \{1_2\}$. Then P is $\Gamma_0(N)$ -conjugate to P' for some number b' if and only if the two fixed points of P are $\Gamma_0(N)$ -conjugate.*

Proof. Assume that the two fixed points of P are $\Gamma_0(N)$ -conjugate. Then there exists an element $T = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \Gamma_0(N)$ such that $\mathfrak{a}, T(\mathfrak{a})$ are the two fixed points of P . Let $P' = T^{-1}PT$. Then $P'(\mathfrak{a}) = \mathfrak{a}$ and $\Gamma_{P'} = \{1_2\}$. Write

$$P' = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a' & b' \\ 0 & d' \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

with $a', d' \in \mathbb{Z}^+$, $a'd' = n$, and $0 \leq b' < |a' - d'|$. Since $P' = T^{-1}PT$, we have

$$(\sigma_{\mathfrak{a}}^{-1} T \sigma_{\mathfrak{a}}) \begin{pmatrix} a' & b' \\ 0 & d' \end{pmatrix} = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} (\sigma_{\mathfrak{a}}^{-1} T \sigma_{\mathfrak{a}}).$$

From this identity, we can deduce that $a' = d$, and hence $d' = a$.

Conversely, suppose that there exists an element $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_0(N)$ such that $T^{-1}PT = P'$, that is,

$$\sigma_{\mathfrak{a}}^{-1} T^{-1} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} = \begin{pmatrix} d & b' \\ 0 & a \end{pmatrix} \sigma_{\mathfrak{a}}^{-1} T^{-1} \sigma_{\mathfrak{a}}.$$

It follows that

$$\left((a-d) \frac{w}{u} - b[w^2, N] \right) u(A + B \frac{w}{u}) = b[w^2, N] \frac{u}{w} (-Cu - Dw).$$

This identity can be written as

$$T(\mathfrak{a}) = \frac{\mathfrak{a}}{1 - \frac{a-d}{b[w^2, N]\mathfrak{a}}},$$

which is the second fixed point of P . That is, the two fixed points of P are $\Gamma_0(N)$ -conjugate. $\square$

Lemma 3.7. *Let $a, d \in \mathbb{Z}^+$, $a \neq d$, $ad = n$, and let $\mathfrak{a} = u/w$ be given as in (3.1) with $(w, N/w) | (a-d)$. Assume that $\mathfrak{a}$ and $\mathfrak{a}' = u_1/w_1$ are the two distinct fixed point of $P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$. Then $(w', N/w') | (a-d)$ for $w' = (w_1, N)$, and*

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}'} \begin{pmatrix} d & b' \\ 0 & a \end{pmatrix} \sigma_{\mathfrak{a}'}^{-1}$$

for some number b' .

Proof. There exist positive integers a', d' with $a'd' = n$ such that

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}'} \begin{pmatrix} a' & b' \\ 0 & d' \end{pmatrix} \sigma_{\mathfrak{a}'}^{-1}$$

for some number b' , and hence we have

$$(3.5) \quad \sigma_{\mathfrak{a}}^{-1} \sigma_{\mathfrak{a}'} \begin{pmatrix} a' & b' \\ 0 & d' \end{pmatrix} = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1} \sigma_{\mathfrak{a}'}.$$

Write $\sigma_{\mathfrak{a}}^{-1} \sigma_{\mathfrak{a}'} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. Then by (3.5) we get that $(d - a')C = 0$, $(a' - a)A = bC$ and $(d - d')D = b'C$. Since $\mathfrak{a} \neq \mathfrak{a}'$, by using (3.3) we find that $C \neq 0$, and hence we have $a' = d$. It follows that $d' = a$. By the argument made in the first paragraph of the proof of Lemma 3.4, we have $(w', N/w')|(a - d)$. $\square$

For a large positive number Y , define

$$D_Y = \{z \in D : \Im \sigma_{\mathfrak{a}_i}^{-1} z < Y, i = 1, 2, \dots, h\}.$$

Let

$$(D_P)_Y = \bigcup_{\gamma \in \Gamma_0(N)} \gamma D_Y.$$

Write

$$c(P)_Y = \int_{(D_P)_Y} k(z, Pz) dz.$$

Lemma 3.8. *Let $\mathfrak{a} = u/w$ be given as in (3.1), and let $P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$ with $a, d \in \mathbb{Z}^+$, $ad = n$, $a \neq d$ be a hyperbolic element of Γ^* with $\Gamma_P = \{1_2\}$ and $P(\mathfrak{a}) = \mathfrak{a}$. Then*

$$c(P)_Y = \frac{\sqrt{n} \ln\{(a - d)^2 [w^2, N] Y / 2\rho\}}{|a - d|} g(\ln \frac{a}{d}) + \int_1^\infty k\left(\frac{(a - d)^2}{n} t\right) \frac{\ln t}{\sqrt{t - 1}} dt + o(1)$$

where $o(1) \rightarrow 0$ as $Y \rightarrow \infty$ and $\rho = C^2 / 2 [C^2 / \ell^2, N] Y$ with $C = (a - d) \frac{w}{u} - b[w^2, N]$ and $\ell = \gcd(C, b[w^2, N] \frac{u}{w})$.

Proof. Let

$$\gamma = \begin{pmatrix} p & r \\ q & s \end{pmatrix}$$

be an element of $SL_2(\mathbb{R})$. The linear fractional transformation, which takes every complex z in the upper half-plane into $\gamma(z)$, maps the horizontal line $\Im z = Y$ into a circle of radius $\frac{1}{2q^2 Y}$ with center at $\frac{p}{q} + \frac{i}{2q^2 Y}$. Let

$$\mu = \begin{pmatrix} 1 & \frac{b}{a-d} \\ 0 & 1 \end{pmatrix}.$$

Then

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \mu^{-1} \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \mu \sigma_{\mathfrak{a}}^{-1}.$$

Note that

$$\mu \sigma_{\mathfrak{a}}^{-1} = \begin{pmatrix} \frac{w/u}{\sqrt{[w^2, N]}} - \frac{b\sqrt{[w^2, N]}}{a-d} & \frac{bu\sqrt{[w^2, N]}}{(a-d)w} \\ -\sqrt{[w^2, N]} & \frac{u}{w}\sqrt{[w^2, N]} \end{pmatrix}.$$

Since

$$P = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \mu^{-1} \begin{pmatrix} 1/y & 0 \\ 0 & y \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \begin{pmatrix} y & 0 \\ 0 & 1/y \end{pmatrix} \mu \sigma_{\mathfrak{a}}^{-1}$$

for any $y \neq 0$, by choosing $y = (a-d)\sqrt{[w^2, N]}$ we obtain that

$$c(P)_Y = \int_{\mu_{\mathfrak{a}}\{(D_P)_Y\}} k(z, \frac{a}{d}z) dz$$

where

$$\mu_{\mathfrak{a}} = \begin{pmatrix} (a-d)\frac{w}{u} - b[w^2, N] & b[w^2, N]\frac{u}{w} \\ \frac{-1}{a-d} & \frac{u}{(a-d)w} \end{pmatrix}.$$

The linear transformation $z \rightarrow \sigma_{\mathfrak{a}} z$ maps the half-plane $\Im z > Y$ into a disk $D_{\mathfrak{a}}$ of radius $\frac{1}{2[w^2, N]Y}$ with center at $\mathfrak{a} + \frac{i}{2[w^2, N]Y}$. Then the transformation $z \rightarrow \mu_{\mathfrak{a}}(z)$ maps the disk $D_{\mathfrak{a}}$ into the half-plane $\Im z > (a-d)^2[w^2, N]Y$. Let

$$\mathfrak{p} = \frac{-b[w^2, N]u/w}{(a-d)\frac{w}{u} - b[w^2, N]}.$$

If

$$\sigma_{\mathfrak{p}} = \begin{pmatrix} \mathfrak{p}\sqrt{[C^2/\ell^2, N]} & 0 \\ \sqrt{[C^2/\ell^2, N]} & 1/\mathfrak{p}\sqrt{[C^2/\ell^2, N]} \end{pmatrix},$$

then $\sigma_{\mathfrak{p}}\infty = \mathfrak{p}$ and $\sigma_{\mathfrak{p}}^{-1}\Gamma_{\mathfrak{p}}\sigma_{\mathfrak{p}} = \Gamma_{\infty}$. By the definition of D_Y , the image of the half-plane $\Im z > Y$ under the linear transformation $z \rightarrow \sigma_{\mathfrak{p}} z$ is not contained in $(D_P)_Y$. Since the linear transformation $z \rightarrow (\mu_{\mathfrak{a}}\sigma_{\mathfrak{p}})(z)$ maps the half-plane $\Im z > Y$ into a disk D_{ρ} of radius ρ centered at $i\rho$, where $\rho = C^2/2[C^2/\ell^2, N]Y$, the disk D_{ρ} is not contained in $\mu_{\mathfrak{a}}\{(D_P)_Y\}$. It follows that

$$\begin{aligned} c(P)_Y &= \int_0^{\pi} d\theta \int_{2\rho \sin \theta}^{(a-d)^2[w^2, N]Y/\sin \theta} k\left(\frac{(a-d)^2}{n \sin^2 \theta}\right) \frac{dr}{r \sin^2 \theta} + o(1) \\ &= \int_1^{\infty} k\left(\frac{(a-d)^2}{n}t\right) \frac{\ln\{(a-d)^2[w^2, N]Yt/2\rho\}}{\sqrt{t-1}} dt + o(1) \end{aligned}$$

where $o(1)$ has a limit zero when $Y \rightarrow \infty$, and hence we have

$$c(P)_Y = \frac{\sqrt{n} \ln\{(a-d)^2[w^2, N]Y/2\rho\}}{|a-d|} g(\ln \frac{a}{d}) + \int_1^{\infty} k\left(\frac{(a-d)^2}{n}t\right) \frac{\ln t}{\sqrt{t-1}} dt + o(1).$$

This completes the proof of the lemma. $\square$

Theorem 3.9. *Let N be any positive integer, and let*

$$H = \{(\mathfrak{a}; a, b, d) : \mathfrak{a}, a, b, \text{ and } d \text{ are given as in Lemma 3.4}\}.$$

Then we have

$$\begin{aligned} \sum_{\{P\}, \Gamma_P = \{1_2\}} c(P)_Y &= \sqrt{n} \ln Y \sum_{\substack{\frac{u}{w}, ad=n, d>0, a \neq d \\ (w, N/w) | (a-d)}} g\left(\ln \frac{a}{d}\right) \\ &+ \frac{\sqrt{n}}{2} \sum_{(\mathfrak{a}; a, b, d) \in H} \ln\{(a-d)^2[w^2, N][\frac{C^2}{\ell^2}, N]/C^2\} g\left(\ln \frac{a}{d}\right) \\ &+ \frac{1}{2} \sum_{\substack{\frac{u}{w}, ad=n, d>0, a \neq d \\ (w, N/w) | (a-d)}} |a-d| \int_1^\infty k\left(\frac{(a-d)^2}{n}t\right) \frac{\ln t}{\sqrt{t-1}} dt + o(1) \end{aligned}$$

where $C = (a-d)\frac{w}{u} - b[w^2, N]$ and $\ell = \gcd(C, b[w^2, N]\frac{u}{w})$.

Proof. Let S_1 be the set of conjugacy classes of hyperbolic elements P of Γ^* such that $\Gamma_P = \{1_2\}$ and such that the two distinct cusps of P are not $\Gamma_0(N)$ -equivalent, and let S_2 be the set of conjugacy classes of hyperbolic elements P of Γ^* such that $\Gamma_P = \{1_2\}$ and such that the two distinct cusps of P are $\Gamma_0(N)$ -equivalent. For the convenience of writing, we denote

$$P(\mathfrak{a}; a, b, d) = \frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}$$

where $\mathfrak{a} = u/w$ is given as in (3.1). Since the two distinct cusps of every element P in S_1 are equivalent to $\Gamma_0(N)$ -inequivalent cusps of the form (3.1), we have

$$\sum_{\{P\} \in S_1} c(P)_Y = \frac{1}{2} \sum_{\substack{(\mathfrak{a}; a, b, d) \in H \\ P(\mathfrak{a}; a, b, d) \in S_1}} c(P(\mathfrak{a}; a, b, d))_Y$$

by Lemma 3.6. It follows from Lemma 3.6 and Lemma 3.7 that

$$\sum_{\{P\} \in S_2} c(P)_Y = \frac{1}{2} \sum_{\substack{(\mathfrak{a}; a, b, d) \in H \\ P(\mathfrak{a}; a, b, d) \in S_2}} c(P(\mathfrak{a}; a, b, d))_Y.$$

It follows that

$$\sum_{(\mathfrak{a}; a, b, d) \in H} c(P(\mathfrak{a}; a, b, d))_Y = 2 \sum_{\{P\}, \Gamma_P = \{1_2\}} c(P)_Y.$$

By Lemma 3.5 and Lemma 3.8, we have

$$\begin{aligned}
& \sum_{(\mathfrak{a}; a, b, d) \in H} c(P(\mathfrak{a}; a, b, d))_Y = 2\sqrt{n} \ln Y \sum_{\substack{\frac{u}{w}, ad=n, d>0, a \neq d \\ (w, N/w) | (a-d)}} g(\ln \frac{a}{d}) \\
& + \sqrt{n} \sum_{(\mathfrak{a}; a, b, d) \in H} \ln \{ (a-d)^2 [w^2, N] [\frac{C^2}{\ell^2}, N] / C^2 \} g(\ln \frac{a}{d}) \\
& + \sum_{\substack{\frac{u}{w}, ad=n, d>0, a \neq d \\ (w, N/w) | (a-d)}} |a-d| \int_1^\infty k \left(\frac{(a-d)^2}{n} t \right) \frac{\ln t}{\sqrt{t-1}} dt + o(1). \quad \square
\end{aligned}$$

3.4. Parabolic components.

Let S be a parabolic element of Γ^* . Then every element of $\Gamma_0(N)$ having the same fixed point as S commutes with S . If $\mathfrak{a} = u/w$ is the fixed point of S , then $\Gamma_S = \Gamma_{\mathfrak{a}}$, and hence we have $\sigma_{\mathfrak{a}}^{-1} \Gamma_S \sigma_{\mathfrak{a}} = \Gamma_\infty$, where $\sigma_{\mathfrak{a}}$ is given as in (3.3). Since $\sigma_{\mathfrak{a}}^{-1} S \sigma_{\mathfrak{a}}$ commutes with every element of Γ_∞ , we have

$$\sigma_{\mathfrak{a}}^{-1} S \sigma_{\mathfrak{a}} = \frac{1}{\sqrt{n}} \begin{pmatrix} a & b \\ 0 & a \end{pmatrix}$$

for some numbers a, b with $a^2 = n$, $a \in \mathbb{Z}$. This implies that Γ^* has parabolic elements only if n is the square of an integer. Furthermore, by (3.4) we see that elements of the form

$$S = \sigma_{\mathfrak{a}} \begin{pmatrix} 1 & b/\sqrt{n} \\ 0 & 1 \end{pmatrix} \sigma_{\mathfrak{a}}^{-1}, \quad 0 \neq b \in \mathbb{Z}$$

constitute a complete set of representatives for the conjugacy classes of parabolic elements of Γ^* having $\mathfrak{a}$ as its fixed point. It follows that

$$\sum_{\{S\}} \int_{D_Y} k(z, Sz) dz = \int_0^Y \int_0^1 \sum_{0 \neq b \in \mathbb{Z}} k(z, z + \frac{b}{\sqrt{n}}) dz + o(1),$$

where the summation on $\{S\}$ is taken over all parabolic classes represented by parabolic elements whose fixed point is $\mathfrak{a} = u/w$ and where $o(1)$ tends to zero as $Y \rightarrow \infty$. Define δ_n to be one if n is the square of an integer and to be zero otherwise.

The identity

$$(3.6) \quad \sigma_{\mathfrak{a}_i} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \sigma_{\mathfrak{a}_j}^{-1} = \begin{pmatrix} A \frac{\mathfrak{a}_i}{\mathfrak{a}_j} - B \mathfrak{a}_i [w^2, N] & B \mathfrak{a}_i \mathfrak{a}_j [w^2, N] \\ \frac{A}{\mathfrak{a}_j} - \frac{D}{\mathfrak{a}_i} - B [w^2, N] & \mathfrak{a}_j B [w^2, N] + D \frac{\mathfrak{a}_j}{\mathfrak{a}_i} \end{pmatrix}$$

holds for $\mathfrak{a}_i = u_i/w$ and $\mathfrak{a}_j = u_j/w$.

Lemma 3.10. *If*

$$\frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$$

is an element of $\sigma_{\mathfrak{a}_i}^{-1} \Gamma^ \sigma_{\mathfrak{a}_i}$, then A, D are integers satisfying $(w, \frac{N}{w}) | (A - D)$. Conversely, if $(w, \frac{N}{w}) | (A - D)$, then there are exactly D elements in $\Gamma_{\mathfrak{a}_i} \setminus \Gamma^*$ represented by matrices of the form $\frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}_i} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \mathfrak{a}_i^{-1}$.*

Proof. Assume that an element $T = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \Gamma_0(N)$ exists such that

$$\sigma_{\mathfrak{a}_i}^{-1} T \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_i} = \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$$

for some positive integers a, d with $ad = n$. Then

$$B = \frac{(\alpha b + \beta d)w^2}{u_i^2[w^2, N]}.$$

Since

$$\frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}_i} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \sigma_{\mathfrak{a}_i}^{-1}$$

is an element of Γ^* , A, D must be integers. By (3.6) we have that

$$B \mathfrak{a}_i^2[w^2, N] = \frac{B u_i^2[w^2, N]}{w^2}$$

is an integer. Since $(u_i, w) = 1$, it follows that

$$\frac{B[w^2, N]}{w} \equiv 0 \pmod{w}.$$

By (3.6) we also have that N/w divides

$$\frac{A - D}{u_i} - \frac{B[w^2, N]}{w}.$$

It follows that

$$\frac{A - D}{u_i} \equiv 0 \pmod{(w, \frac{N}{w})}.$$

Since $(u_i, w) = 1$, we have that

$$A - D \equiv 0 \pmod{(w, \frac{N}{w})}.$$

Conversely, if $(w, \frac{N}{w}) | (A - D)$, then it follows from the second part of the proof of Lemma 3.4 that

$$\frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}_i} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \sigma_{\mathfrak{a}_i}^{-1}$$

is an element of Γ^* for some number of B . By an argument similar to that made in the proof of the lemma 3.5, there are exactly D elements in $\Gamma_{\mathfrak{a}_i} \setminus \Gamma^*$ represented by matrices of the form $\frac{1}{\sqrt{n}} \sigma_{\mathfrak{a}_i} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \mathfrak{a}_i^{-1}$.

This completes the proof of the lemma. $\square$

The Maass-Selberg relation [7]. Let Γ be a discontinuous group of finite type, and let $\mathfrak{a}_1, \dots, \mathfrak{a}_h$ be a complete set of inequivalent cusps of Γ . Assume that $f(z, s)$ and $g(z, s')$ are two automorphic functions with respect to Γ such that

$$\Delta f(z, s) = s(1-s)f(z, s)$$

and

$$\Delta g(z, s') = s'(1-s')g(z, s')$$

and such that the constant terms of the Fourier expansion at $\mathfrak{a}_j$ of f and g are given respectively by

$$c_j y^s + c'_j y^{1-s}$$

and

$$d_j y^{s'} + d'_j y^{1-s'}.$$

Then, for the compact parts f^Y and g^Y of f and g , we have

$$\begin{aligned} & \langle f^Y(z, s), \bar{g}^Y(z, s') \rangle \\ &= \sum_{j=1}^h \left(\frac{c_j d_j Y^{s+s'-1} - c'_j d'_j Y^{-s-s'+1}}{s+s'-1} + \frac{c_j d'_j Y^{s-s'} - c'_j d_j Y^{-s+s'}}{s-s'} \right) \end{aligned}$$

whenever $s(1-s) \neq s'(1-s')$ and f^Y and g^Y are square integrable on $\Gamma \backslash \mathcal{H}$.

Theorem 3.11. We have

$$\begin{aligned} & \sum_{i=1}^{\nu(N)} \int_{D_Y} \left\{ \sum_{ad=n, 0 \leq b < d} E_i \left(\frac{az+b}{d}, s \right) \right\} E_i(z, \bar{s}) dz \\ &= \sum_{j=1}^{\nu(N)} \sum_{\substack{a, d > 0, ad=n \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, N/w_j) | (a-d)}} a^s d^{1-s} \left\{ \frac{Y^{s+\bar{s}-1} - \sum_{i=1}^{\nu(N)} \varphi_{ij}(s) \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s+\bar{s}-1} \right. \\ & \quad \left. + \frac{\varphi_{jj}(\bar{s}) Y^{s-\bar{s}} - \varphi_{jj}(s) Y^{-s+\bar{s}}}{s-\bar{s}} \right\} - \sum_{j=1}^{\nu(N)} \sum_{a, d, b}^* \frac{\left(\frac{A'}{D'} \right)^{1-s} \varphi_{jj'}(s) Y^{-s+\bar{s}}}{s-\bar{s}} + o_Y(1) \end{aligned}$$

as $\text{Re } s \rightarrow 1/2$, where the sum on a, d, b has the same meaning as in (3.10) with A', D' being defined there and $o_Y(1)$ tends to zero as $Y \rightarrow \infty$.

Proof. Let

$$f_i(z) = \sum_{ad=n, 0 \leq b < d} E_i \left(\frac{az+b}{d}, s \right).$$

Then $f_i(z)$ is an automorphic function with respect to $\Gamma_0(N)$. Let the constant term of the Fourier expansion at $\mathfrak{a}_j$ of $f_i(z)$ be $c_{ij}y^s + c'_{ij}y^{1-s}$. By the Maass-Selberg relation we have

$$(3.7) \quad \begin{aligned} & \sum_{i=1}^{\nu(N)} \int_{D_Y} f_i(z) E_i(z, \bar{s}) dz - o_Y(1) \\ &= \sum_{i,j=1}^{\nu(N)} \left(\frac{c_{ij} \delta_{ij} Y^{s+\bar{s}-1} - c'_{ij} \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s + \bar{s} - 1} + \frac{c_{ij} \varphi_{ij}(\bar{s}) Y^{s-\bar{s}} - c'_{ij} \delta_{ij} Y^{-s+\bar{s}}}{s - \bar{s}} \right), \end{aligned}$$

where $o_Y(1)$ tends to zeros as $Y \rightarrow \infty$.

Since

$$E_i \left(\frac{a\sigma_{\mathfrak{a}_j} z + b}{d}, s \right) = \sum_{\gamma \in \Gamma_{\mathfrak{a}_i} \setminus \Gamma_0(N)} y \left(\sigma_{\mathfrak{a}_i}^{-1} \gamma \frac{a\sigma_{\mathfrak{a}_j} z + b}{d} \right)^s,$$

we have

$$f_i(\sigma_{\mathfrak{a}_j} z) = \sum_{ad=n, 0 \leq b < d} \sum_{\gamma \in \Gamma_{\mathfrak{a}_i} \setminus \Gamma_0(N)} y \left(\sigma_{\mathfrak{a}_i}^{-1} \gamma \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_j} z \right)^s.$$

Let $a, d > 0$ and b be integers with $ad = n$ and $0 \leq b < d$. If the matrix equation

$$(3.8) \quad \sigma_{\mathfrak{a}_j}^{-1} \frac{1}{\sqrt{n}} T \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_j} = \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$$

holds for an element T in $\Gamma_0(N)$ and for some numbers A, D and B , then A, D are integers with $AD = n$. Moreover, by Lemma 3.10 we must have $(w_i, N/w_i) | (A - D)$ for $\mathfrak{a}_i = u_i/w_i$. Note that, for any fixed pair of positive integers A, D with $AD = n$, if the matrix equation (3.8) has a solution T, a, b, d for a number B with $0 \leq B < D$, then the matrix equation (3.8) has solutions for exactly D number of B 's with $0 \leq B < D$ by the proof of Lemma 3.5. By Theorem 2.7 of Iwaniec [6], we have the following decomposition

$$\begin{aligned} & \Gamma_\infty \setminus \sigma_{\mathfrak{a}_i}^{-1} \frac{1}{\sqrt{n}} \Gamma_0(N) \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_j} \\ &= \left[\delta_{ij} \cup \bigcup_{\gamma > 0} \bigcup_{\delta \pmod{\gamma}} \begin{pmatrix} * & * \\ \gamma & \delta \end{pmatrix} \Gamma_\infty \right] \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \end{aligned}$$

where the unions on γ, δ run over all numbers $\gamma > 0, 0 \leq \delta < \gamma$ such that $\begin{pmatrix} * & * \\ \gamma & \delta \end{pmatrix}$ belong to $\sigma_{\mathfrak{a}_i}^{-1} \Gamma_0(N) \sigma_{\mathfrak{a}_j}$. Then it follows from Lemma 3.10 and the proof of Theorem 3.4 of Iwaniec [6] that the constant coefficient in the Fourier expansion at infinity of the function

$$\tilde{f}_i(\sigma_{\mathfrak{a}_j} z) = \sum_{*} \sum_{\gamma \in \Gamma_{\mathfrak{a}_i} \setminus \Gamma_0(N)} y \left(\sigma_{\mathfrak{a}_i}^{-1} \gamma \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_j} z \right)^s,$$

where the sum $\sum_*$ runs over all integers $a, d > 0, 0 \leq b < d, ad = n$ such that the matrix equation (3.8) has a solution T, A, B, D , is equal to

$$(3.9) \quad \sum_{\substack{A, D > 0, AD = n \\ (w_j, N/w_j) | (A - D)}} \{ \delta_{ij} A^s D^{1-s} y^s + \varphi_{ij}(s) A^{1-s} D^s y^{1-s} \}.$$

Next, we assume that $a, d > 0$ and b are integers with $ad = n$ and $0 \leq b < d$ such that the matrix equation (3.8) has no solutions. Then the cusp $(a\mathfrak{a}_j + b)/d$ is not $\Gamma_0(N)$ -equivalent to $\mathfrak{a}_j$. Because every cusp of $\Gamma_0(N)$ is $\Gamma_0(N)$ -equivalent to a cusp given in (3.1), a cusp $\mathfrak{a}_{j'}$, $j' \neq j$, exists which is equivalent to $(a\mathfrak{a}_j + b)/d$ under $\Gamma_0(N)$. That is

$$\mathfrak{a}_{j'} = \gamma_0 \left(\frac{a\mathfrak{a}_j + b}{d} \right)$$

for some element γ_0 in $\Gamma_0(N)$. Let $\sigma_{\mathfrak{a}_{j'}}$ be the element in $PSL_2(\mathbb{R})$ such that $\sigma_{\mathfrak{a}_{j'}} \infty = \mathfrak{a}_{j'}$ and $\sigma_{\mathfrak{a}_{j'}}^{-1} \Gamma_{\mathfrak{a}_{j'}} \sigma_{\mathfrak{a}_{j'}} = \Gamma_\infty$, and let

$$T_0 = \sigma_{\mathfrak{a}_{j'}}^{-1} \gamma_0 \frac{1}{\sqrt{n}} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_j}.$$

Then T_0 is an element of $SL_2(\mathbb{R})$ satisfying $T_0(\infty) = \infty$. It follows that we have the decomposition

$$\Gamma_\infty \backslash \sigma_{\mathfrak{a}_i}^{-1} \frac{1}{\sqrt{n}} \Gamma_0(N) \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \sigma_{\mathfrak{a}_j} = \left[\bigcup_{\gamma > 0} \bigcup_{\delta \pmod{\gamma}} \begin{pmatrix} * & * \\ \gamma & \delta \end{pmatrix} \Gamma_\infty \right] T_0$$

where the unions on γ, δ run over all numbers $\gamma > 0, 0 \leq \delta < \gamma$ such that $\begin{pmatrix} * & * \\ \gamma & \delta \end{pmatrix}$ belong to $\sigma_{\mathfrak{a}_i}^{-1} \Gamma_0(N) \sigma_{\mathfrak{a}_{j'}}$. Note that, since $f_i(\sigma_{\mathfrak{a}_j} z)$ and $\tilde{f}_i(\sigma_{\mathfrak{a}_j} z)$ are both periodic function of $x, z = x + iy$, of period 1, the function

$$\tilde{g}_i(\sigma_{\mathfrak{a}_j} z) = f_i(\sigma_{\mathfrak{a}_j} z) - \tilde{f}_i(\sigma_{\mathfrak{a}_j} z)$$

is also a periodic function of x of period 1. Since $T_0(\infty) = \infty$, we can write

$$T_0 = \frac{1}{\sqrt{n}} \begin{pmatrix} A' & B' \\ 0 & D' \end{pmatrix}$$

for some real numbers A', B' and D' . Then it follows the above decomposition, the above note, and the proof of Theorem 3.4 of Iwaniec [6] that the constant coefficient in the Fourier expansion at infinity of the function $\tilde{g}_i(\sigma_{\mathfrak{a}_j} z)$ is equal to

$$(3.10) \quad \sum_{a, d, b}^* \varphi_{ij'}(s) \left(\frac{A'}{D'} y \right)^{1-s}$$

where the sum is over all integers $a, d > 0$ and $0 \leq b < d$ with $ad = n$ such that the matrix equation (3.8) has no solutions. It follows from (3.7), (3.9) and (3.10) that

$$\begin{aligned}
& \sum_{i=1}^{\nu(N)} \int_{D_Y} f_i(z) E_i(z, \bar{s}) dz - o_Y(1) \\
&= \sum_{i,j=1}^{\nu(N)} \sum_{\substack{A,D>0, AD=n \\ (w_j, N/w_j) | (A-D)}} \left\{ \frac{A^s D^{1-s} \delta_{ij} Y^{s+\bar{s}-1} - \varphi_{ij}(s) A^{1-s} D^s \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s + \bar{s} - 1} \right. \\
&\quad \left. + \frac{\delta_{ij} A^s D^{1-s} \varphi_{ij}(\bar{s}) Y^{s-\bar{s}} - \varphi_{ij}(s) A^{1-s} D^s \delta_{ij} Y^{-s+\bar{s}}}{s - \bar{s}} \right\} \\
&- \sum_{i,j=1}^{\nu(N)} \sum_{a,d,b} \left\{ \frac{\left(\frac{A'}{D'}\right)^{1-s} \varphi_{ij'}(s) \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s + \bar{s} - 1} + \frac{\delta_{ij} \left(\frac{A'}{D'}\right)^{1-s} \varphi_{ij'}(s) Y^{-s+\bar{s}}}{s - \bar{s}} \right\}
\end{aligned}$$

where the sum over a, d, b has the same meaning as in (3.10). If $\Phi(s)$ is the constant term matrix of the Eisenstein series for $\Gamma_0(N)$, then $\Phi(s)$ is a symmetric matrix and satisfies the identity

$$\Phi(s) \Phi(1-s) = I$$

for all complex s , where I is the identity matrix; see Kubota [7]. Since $j \neq j'$, we have

$$\sum_{i=1}^{\nu(N)} \varphi_{ij'}(s) \varphi_{ij}(\bar{s}) = 0$$

as $\text{Re } s \rightarrow 1/2$. Hence, the above identity can be written as

$$\begin{aligned}
& \sum_{i=1}^{\nu(N)} \int_{D_Y} f_i(z) E_i(z, \bar{s}) dz = \sum_{i,j=1}^{\nu(N)} \sum_{\substack{A,D>0, AD=n \\ (w_j, N/w_j) | (A-D)}} A^s D^{1-s} \\
(3.11) \quad & \times \left\{ \frac{\delta_{ij} Y^{s+\bar{s}-1} - \varphi_{ij}(s) \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s + \bar{s} - 1} + \delta_{ij} \frac{\varphi_{ij}(\bar{s}) Y^{s-\bar{s}} - \varphi_{ij}(s) Y^{-s+\bar{s}}}{s - \bar{s}} \right\} \\
& - \sum_{i,j=1}^{\nu(N)} \sum_{a,d,b}^* \frac{\delta_{ij} \left(\frac{A'}{D'}\right)^{1-s} \varphi_{ij'}(s) Y^{-s+\bar{s}}}{s - \bar{s}} + o_Y(1)
\end{aligned}$$

as $\text{Re } s \rightarrow 1/2$, where the sum on a, d, b has the same meaning as in (3.10).

This completes the proof of the theorem. $\square$

Theorem 3.12. *Put*

$$c(\infty)_Y = \delta_n \nu(N) \int_0^Y \int_0^1 \sum_{0 \neq b \in \mathbb{Z}} k(z, z + \frac{b}{\sqrt{n}}) dz - \int_{D_Y} H(z, z) dz.$$

Then

$$\begin{aligned}
\frac{c(\infty)_Y}{\sqrt{n}} &= \delta_n \nu(N) g(0) \ln \frac{\sqrt{n}}{2} - \sum_{j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0, a \neq d \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} g\left(\ln \frac{a}{d}\right) \ln Y \\
&\quad - \frac{\delta_n \nu(N)}{2\pi} \int_{-\infty}^{\infty} h(r) \frac{\Gamma'}{\Gamma} (1+ir) dr + \frac{1}{4} h(0) \{ \delta_n \nu(N) \\
&\quad + \sum_{j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0, a \neq d \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} \varphi_{jj}\left(\frac{1}{2}\right) \} \\
&\quad + \sum_{i,j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0 \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | a-d}} \frac{1}{4\pi} \int_{-\infty}^{\infty} h(r) \left(\frac{a}{d}\right)^{ir} \varphi'_{ij}\left(\frac{1}{2} + ir\right) \varphi_{ij}\left(\frac{1}{2} - ir\right) dr \\
&\quad - \frac{1}{8} h(0) \sum_{j=1}^{\nu(N)} \sum_{a,d,b}^* \left(\frac{A'}{D'}\right)^{1/2} \varphi_{jj'}\left(\frac{1}{2}\right) + o_Y(1),
\end{aligned}$$

where $o_Y(1)$ tends to zero as $Y \rightarrow \infty$, and the sum on a, d, b has the same meaning as in (3.10) with A', D' being defined there.

Proof. By the argument of [7], pp.102–106 we have

$$\begin{aligned}
&\frac{1}{\sqrt{n}} \int_0^Y \int_0^1 \sum_{0 \neq b \in \mathbb{Z}} k\left(z, z + \frac{b}{\sqrt{n}}\right) dz \\
&= g(0) \ln(\sqrt{n}Y) - \frac{1}{2\pi} \int_{-\infty}^{\infty} h(r) \frac{\Gamma'}{\Gamma} (1+ir) dr - g(0) \ln 2 + \frac{1}{4} h(0) + o(1).
\end{aligned}$$

Let

$$\varphi_{ij,m}(s) = \sum_c \frac{1}{|c|^{2s}} \left(\sum_d e\left(m \frac{d}{c}\right) \right)$$

where the sum runs over $c > 0, d$ modulo c with $\begin{pmatrix} * & * \\ c & d \end{pmatrix} \in \sigma_{\mathfrak{a}_i}^{-1} \Gamma_0(N) \sigma_{\mathfrak{a}_j}$. Then we have

$$\begin{aligned}
E_i(\sigma_{\mathfrak{a}_j} z, s) &= \delta_{ij} y^s + \varphi_{ij}(s) y^{1-s} \\
&\quad + \frac{2\pi^s \sqrt{y}}{\Gamma(s)} \sum_{m \neq 0} |m|^{s-\frac{1}{2}} K_{s-\frac{1}{2}}(2|m|\pi y) \varphi_{ij,m}(s) e(mx)
\end{aligned}$$

where

$$\varphi_{ij}(s) = \frac{\sqrt{\pi} \Gamma(s - \frac{1}{2})}{\Gamma(s)} \varphi_{ij,0}(s).$$

By Theorem 3.11, we have

$$\begin{aligned}
& \sum_{i=1}^{\nu(N)} \int_{D_Y} \left\{ \sum_{ad=n, 0 \leq b < d} E_i \left(\frac{az+b}{d}, s \right) \right\} E_i(z, \bar{s}) dz \\
&= \sum_{i,j=1}^{\nu(N)} \sum_{\substack{a,d>0, ad=n \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} a^s d^{1-s} \left\{ \frac{\delta_{ij} Y^{s+\bar{s}-1} - \varphi_{ij}(s) \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s + \bar{s} - 1} \right. \\
&\quad \left. + \delta_{ij} \frac{\varphi_{ij}(\bar{s}) Y^{s-\bar{s}} - \varphi_{ij}(s) Y^{-s+\bar{s}}}{s - \bar{s}} \right\} - \sum_{i,j=1}^{\nu(N)} \sum_{a,d,b}^* \frac{\delta_{ij} \left(\frac{A'}{D'} \right)^{1-s} \varphi_{ij'}(s) Y^{-s+\bar{s}}}{s - \bar{s}} + o_Y(1)
\end{aligned}$$

as $\text{Re } s \rightarrow 1/2$, and hence we have

$$\begin{aligned}
& \sum_{i=1}^{\nu(N)} \int_{D_Y} \left\{ \sum_{ad=n, 0 \leq b < d} E_i \left(\frac{az+b}{d}, s \right) \right\} E_i(z, \bar{s}) dz \\
&= \sum_{j=1}^{\nu(N)} \sum_{\substack{a,d>0, ad=n \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} a^s d^{1-s} \left\{ \frac{Y^{s+\bar{s}-1} - \sum_{i=1}^{\nu(N)} \varphi_{ij}(s) \varphi_{ij}(\bar{s}) Y^{-s-\bar{s}+1}}{s + \bar{s} - 1} \right. \\
&\quad \left. + \frac{\varphi_{jj}(\bar{s}) Y^{s-\bar{s}} - \varphi_{jj}(s) Y^{-s+\bar{s}}}{s - \bar{s}} \right\} - \sum_{j=1}^{\nu(N)} \sum_{a,d,b}^* \frac{\left(\frac{A'}{D'} \right)^{1-s} \varphi_{jj'}(s) Y^{-s+\bar{s}}}{s - \bar{s}} + o_Y(1)
\end{aligned}$$

as $\text{Re } s \rightarrow 1/2$, where the sum on u/w runs over all cusps given in (3.1) and the sum on a, d, b has the same meaning as in (3.10) with A', D' being defined there and $o_Y(1)$ tends to zero as $Y \rightarrow \infty$.

By partial integration, we obtain

$$h(r) = \frac{1}{r^4} \int_0^\infty g^{(4)}(\ln u) u^{ir-1} du$$

for nonzero r . Then it follows that

$$\begin{aligned}
 (3.12) \quad & \lim_{S \rightarrow \frac{1}{2}^+} \int_{-\infty}^{\infty} h(r) \left\{ \sum_{i=1}^{\nu(N)} \int_{D_Y} \left(\sum_{ad=n, 0 \leq b < d} E_i\left(\frac{az+b}{d}, S+ir\right) E_i(z, S-ir) dz \right) \right\} dr \\
 &= \sqrt{n} \sum_{j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0 \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | a-d}} \left\{ 4\pi g\left(\ln \frac{a}{d}\right) \ln Y - \sum_{i=1}^{\nu(N)} \int_{-\infty}^{\infty} h(r) \left(\frac{a}{d}\right)^{ir} \varphi'_{ij}\left(\frac{1}{2} + ir\right) \right. \\
 &\quad \times \varphi_{ij}\left(\frac{1}{2} - ir\right) dr + \int_{-\infty}^{\infty} h(r) \left(\frac{a}{d}\right)^{ir} \frac{\varphi_{jj}\left(\frac{1}{2} - ir\right) Y^{2ir}}{ir} dr \Big\} \\
 &\quad - \sum_{j=1}^{\nu(N)} \sum_{a,d,b}^* \int_{-\infty}^{\infty} h(r) \varphi_{jj'}\left(\frac{1}{2} + ir\right) \left(\frac{A'}{D'}\right)^{\frac{1}{2}-ir} \frac{Y^{-2ir}}{2ir} dr + o_Y(1).
 \end{aligned}$$

By the Riemann-Lebesgue theorem (cf. §1.8 of [16]), we have

$$\begin{aligned}
 (3.13) \quad & \lim_{Y \rightarrow \infty} \int_{-\infty}^{\infty} h(r) \left(\frac{a}{d}\right)^{ir} \frac{\varphi_{jj}\left(\frac{1}{2} - ir\right) Y^{2ir}}{ir} dr \\
 &= \lim_{Y \rightarrow \infty} \int_{-\infty}^{\infty} h(r) \Re \left(\varphi_{jj}\left(\frac{1}{2} - ir\right) \right) \frac{\sin\{r \ln(aY^2/d)\}}{r} dr = \pi h(0) \varphi_{jj}\left(\frac{1}{2}\right).
 \end{aligned}$$

By (3.12) and (3.13), we have

$$\begin{aligned}
 & \frac{1}{\sqrt{n}} \int_{D_Y} H(z, z) dz = \sum_{j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0 \\ \mathfrak{a}_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | a-d}} \left\{ g\left(\ln \frac{a}{d}\right) \ln Y + \frac{1}{4} h(0) \varphi_{jj}\left(\frac{1}{2}\right) \right. \\
 & \quad \left. - \sum_{i=1}^{\nu(N)} \frac{1}{4\pi} \int_{-\infty}^{\infty} h(r) \left(\frac{a}{d}\right)^{ir} \varphi'_{ij}\left(\frac{1}{2} + ir\right) \varphi_{ij}\left(\frac{1}{2} - ir\right) dr \right\} \\
 & \quad + \frac{1}{8} h(0) \sum_{j=1}^{\nu(N)} \sum_{a,d,b}^* \left(\frac{A'}{D'}\right)^{1/2} \varphi_{jj'}\left(\frac{1}{2}\right) + o_Y(1).
 \end{aligned}$$

The stated identity then follows. $\square$

It follows from Theorem 3.9 and Theorem 3.12 that
(3.14)

$$\begin{aligned}
& \lim_{Y \rightarrow \infty} \left(c(\infty)_Y + \sum_{\{P\}, \Gamma_P = \{1_2\}} c(P)_Y \right) \\
&= \sqrt{n} \delta_n \nu(N) g(0) \ln \frac{\sqrt{n}}{2} + \frac{\sqrt{n}}{4} h(0) \{ \delta_n \nu(N) + \sum_{j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0, a \neq d \\ a_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} \varphi_{jj}(\frac{1}{2}) \} \\
&+ \frac{\sqrt{n}}{2} \sum_{w|N, w>0} \sum_{ad=n, d>0, a \neq d} \sum_b \frac{\ln \{ (a-d)^2 w N [\frac{C^2}{\ell^2}, N] / C^2 \}}{|a-d|} g(\ln \frac{a}{d}) \\
&- \frac{\delta_n \nu(N) \sqrt{n}}{2\pi} \int_{-\infty}^{\infty} h(r) \frac{\Gamma'}{\Gamma} (1+ir) dr \\
&+ \sum_{j=1}^{\nu(N)} \sum_{\substack{ad=n, d>0 \\ a_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} \frac{\sqrt{n}}{4\pi} \int_{-\infty}^{\infty} h(r) \left(\frac{a}{d} \right)^{ir} \sum_{i=1}^{\nu(N)} \varphi'_{ij}(\frac{1}{2} + ir) \varphi_{ij}(\frac{1}{2} - ir) dr \\
&+ \sum_{\substack{ad=n, d>0, a \neq d \\ a_j = \frac{u_j}{w_j}, (w_j, \frac{N}{w_j}) | (a-d)}} \frac{1}{2} |a-d| \int_1^{\infty} k \left(\frac{(a-d)^2}{n} t \right) \frac{\ln t}{\sqrt{t-1}} dt \\
&- \frac{1}{8} h(0) \sum_{j=1}^{\nu(N)} \sum_{a, d, b}^* \left(\frac{A'}{D'} \right)^{1/2} \varphi_{jj'}(\frac{1}{2})
\end{aligned}$$

with $C = (a-d)w - bwN$, where $\ell = (C, bN)$ and the summation on b is taken over all numbers b such that $C, bN/w \in \mathbb{Z}$, $N|C$ and $0 \leq b < |a-d|$. Note that there are exactly $|a-d|$ number of such numbers b by Lemma 3.5.

Denote by $c(\infty)$ the right side of the identity (3.14). We conclude that the trace formula (2.3) can be written as

$$\begin{aligned}
& d(n) h(-\frac{i}{2}) + \sqrt{n} \sum_{j=1}^{\infty} h(\kappa_j) \text{tr}_j T_n \\
&= c(I) + \sum_{\{R\}} c(R) + \sum_{\{P\}, \Gamma_P \neq \{1_2\}} c(P) + c(\infty)
\end{aligned}
\tag{3.15}$$

for $\text{Re } s > 1$, where the summations on the right side of the identity are taken over the conjugacy classes.

Theorem 3.13. *Let $c(P) = \int_{\Gamma_P \backslash \mathcal{H}} k(z, Pz) dz$ for hyperbolic elements $P \in \Gamma_0(N)$. Then the series*

$$\sum_{\{P\}} c(P)$$

represents an analytic function in the half-plane $\text{Re } s > 0$ except for having a possible pole at $s = 1/2$ and for having simple poles at $s = 1, \frac{1}{2} \pm i\kappa_j$, $j = 1, 2, \dots$.

Proof. We have

$$g^{(4)}(\log u) = A(s)u^{\frac{1}{2}-s} + O_s(u^{-\frac{1}{2}}),$$

where $A(s)$ is an analytic function of s for $\text{Re } s > 0$ and, for every complex number s with $\text{Re } s > 0$, there exists a finite constant $B(s)$ depending only on s such that

$$|O_s(u^{-\frac{1}{2}})| \leq B(s)u^{-\frac{1}{2}}.$$

Moreover, for every fixed value of u , the term $O_s(u^{-\frac{1}{2}})$ also represents an analytic function of s for $\text{Re } s > 0$. Since

$$h(r) = \frac{1}{r^4} \int_0^\infty g^{(4)}(\ln u) u^{ir-1} du$$

for nonzero r , we have

$$h(r) = \frac{A(s)}{r^4} \left(\frac{1}{s - \frac{1}{2} - ir} + \frac{1}{s - \frac{1}{2} + ir} \right) + O_s(r^{-4})$$

for $\text{Re } s > 1$ and for nonzero r with $|\text{Im } r| < \frac{1}{2} - \epsilon$. By analytic continuation, we obtain that

$$(3.16) \quad h(r) = \frac{2A(s)(s - \frac{1}{2})}{r^4[(s - \frac{1}{2})^2 + r^2]} + O_s(r^{-4})$$

for $\text{Re } s > 0$ and for nonzero r with $|\text{Im } r| < \frac{1}{2} - \epsilon$. It follows from results of [17] and (2.3) that the left side of (3.15) is an analytic function of s for $\text{Re } s > 0$ except for having simple poles at $s = 1, \frac{1}{2} \pm i\kappa_j$, $j = 1, 2, \dots$. Then the right side of (3.15) can be interpreted as an analytic function of s in the same region by analytic continuation.

Since $k(t) = (1+t/4)^{-s}$, by Lemma 3.1 we have that $c(R)$ is analytic for $\text{Re } s > 0$ except for a simple pole at $s = 1/2$. There are only a finite number of elliptic conjugacy classes $\{R\}$. The term $c(I)$ is a constant.

Since $g(0) = 2\sqrt{\pi}\Gamma(s - \frac{1}{2})\Gamma(s)^{-1}$,

$$h(0) = 2\sqrt{\pi}4^s \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \int_1^\infty (u + \frac{1}{u} + 2)^{\frac{1}{2}-s} \frac{du}{u}$$

and

$$g(\ln \frac{a}{d}) = 2\sqrt{\pi}4^{s-\frac{1}{2}} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \left(\frac{a}{d} + \frac{d}{a} + 2 \right)^{\frac{1}{2}-s},$$

the sum of first three terms and the last two terms on the right side of the identity (3.14) is analytic for $\operatorname{Re} s > 0$ except for a pole at $s = 1/2$.

By Stirling's formula the identity

$$(3.17) \quad \frac{\Gamma'(z)}{\Gamma(z)} = \ln z + O(1)$$

holds uniformly when $|\arg z| \leq \pi - \delta$ for a small positive number δ . It follows from (3.13) and (3.17) that the fourth term on the right side of the identity (3.14) is analytic for $\operatorname{Re} s > 0$ except for a possible pole at $s = 1/2$.

By Theorem 4.4.1 of Kubota [7], each Eisenstein series $E_i(z, s)$ has a meromorphic continuation to the whole s -plane, and the identity

$$\sum_{i=1}^{\nu(N)} \varphi_{ij} \left(\frac{1}{2} + ir \right) \varphi_{ij} \left(\frac{1}{2} - ir \right) = 1$$

hold for all real r and for $i = 1, 2, \dots, \nu(N)$. It follows that functions $\varphi_{ij}(s)$, $i, j = 1, 2, \dots, \nu(N)$, are analytic on the line $\operatorname{Re} s = 1/2$. Let Y be a fixed large positive number. By the Maass-Selberg relation, we have

$$(3.18) \quad \begin{aligned} & \operatorname{Re} \sum_{i=1}^{\nu(N)} \varphi'_{ij} \left(\frac{1}{2} + ir \right) \varphi_{ij} \left(\frac{1}{2} - ir \right) \\ &= 2 \ln Y + \frac{\varphi_{jj}(1/2 - ir)Y^{2ir} - \varphi_{jj}(1/2 + ir)Y^{-2ir}}{2ir} \\ & \quad - \int_{D_Y} E_j \left(z, \frac{1}{2} + ir \right) E_j \left(z, \frac{1}{2} - ir \right) dz + o(1). \end{aligned}$$

Let $\mathfrak{a}_i = u_i/w_i$ be a cusp given in (3.1), and let $\eta = 1/2 + ir$. By the argument following Lemma 3.6 of [3], we have

$$(3.19) \quad \begin{aligned} E_i(z, \eta) &= \delta_{\mathfrak{a}_i \infty} y^\eta + \frac{\sqrt{\pi} \Gamma(\eta - \frac{1}{2})}{\Gamma(\eta)} \varphi_{\mathfrak{a}_i \infty, 0}(\eta) y^{1-\eta} \\ & \quad + \frac{2\pi^\eta \sqrt{y}}{\Gamma(\eta)} \sum_{m \neq 0} |m|^{\eta - \frac{1}{2}} K_{\eta - \frac{1}{2}}(2|m|\pi y) \varphi_{\mathfrak{a}_i \infty, m}(\eta) e(mx) \end{aligned}$$

where

$$\begin{aligned} & \varphi_{\mathfrak{a}_i \infty, 0}(\eta) \\ &= \frac{\varphi(w_i)}{\varphi((w_i, N/w_i))} \left(\frac{(w_i, N/w_i)}{w_i N} \right)^\eta \prod_{p|N} \frac{1}{1 - p^{-2\eta}} \prod_{p|\frac{N}{w_i}} (1 - p^{1-2\eta}) \frac{\zeta(2\eta - 1)}{\zeta(2\eta)} \end{aligned}$$

and

$$\varphi_{\mathfrak{a}_i\infty,m}(\eta) = \left(\frac{(w_i, N/w_i)}{w_i N} \right)^\eta \sum_{(c, N/w_i)=1} \frac{1}{c^{2\eta}} \sum_{\substack{d \bmod(cw_i), (d, cw_i)=1 \\ cd \equiv u_i \bmod(w_i, N/w_i)}} e\left(-\frac{md}{cw_i}\right).$$

It follows from the functional identity of the Riemann zeta-function that

$$\begin{aligned} \frac{\Gamma(\eta - \frac{1}{2})}{\Gamma(\eta)} \varphi_{\mathfrak{a}_i\infty,0}(\eta) &= \pi^{2\eta - \frac{3}{2}} \frac{\Gamma(1-\eta)}{\Gamma(\eta)} \frac{\zeta(2-2\eta)}{\zeta(2\eta)} \frac{\varphi(w_i)}{\varphi((w_i, N/w_i))} \\ &\quad \times \left(\frac{(w_i, N/w_i)}{w_i N} \right)^\eta \prod_{p|N} \frac{1}{1-p^{-2\eta}} \prod_{p|\frac{N}{w_i}} (1-p^{1-2\eta}). \end{aligned}$$

By using Stirling's formula

$$|\Gamma(\sigma + it)| \sim \sqrt{2\pi} e^{-\pi|t|/2} |t|^{\sigma-1/2}$$

for any fixed real value of σ as $t \rightarrow \infty$, we obtain that

$$(3.20) \quad \frac{\Gamma(\eta - \frac{1}{2})}{\Gamma(\eta)} \varphi_{\mathfrak{a}_i\infty,0}(\eta) y^{1-\eta} \ll \sqrt{y} \ln^2(|r| + 1).$$

By (1.1) and by partial integration, we find that

$$(3.21) \quad \frac{K_{\eta-1/2}(2|m|\pi y)}{\Gamma(\eta)} = \frac{2^{\eta-1/2}}{(2|m|\pi y)^{\eta-1/2} \sqrt{\pi}} \int_0^\infty \frac{\cos(2|m|\pi y t)}{(1+t^2)^\eta} dt \ll \frac{1+|r|^3}{|m|^3 y^3}.$$

Then it follows from (3.16), (3.18)-(3.21), Lemma 4.7 of [3] together with the proof of Theorem 1 in [8], and Cauchy's inequality that the fifth term on the right side of the identity (3.14) is analytic for $\operatorname{Re} s > 0$ except for a possible pole at $s = 1/2$.

Also, it follows from (3.12) and Lemma 8.2 of Strömbergsson [15] that the fifth term on the right side of the identity (3.14) is analytic for $\operatorname{Re} s > 0$ except for a possible pole at $s = 1/2$.

Therefore, by (3.15) we have proved that the series

$$\sum_{\{P\}} c(P)$$

represents an analytic function of s in the half-plane $\operatorname{Re} s > 0$ except for having a possible pole at $s = 1/2$ and for having simple poles at $s = 1, \frac{1}{2} \pm i\kappa_j$, $j = 1, 2, \dots$. $\square$

4. PROOF OF THE MAIN THEOREM

Proofs for the following lemmas from Lemma 4.1 to Lemma 4.4 are minor modifications of corresponding lemmas in Conrey-Li [2]. For the convenience of readers, we sketch the proofs here.

Lemma 4.1. *Let $\lambda > 0$ be an eigenvalue of Δ for $\Gamma_0(N)$ with N square free. Assume that $(n, N) = 1$. Put $\tau = 1/2 + i\kappa$ with $\kappa = \sqrt{\lambda - 1/4}$. Then we have*

$$4^\tau \sqrt{\pi n} \frac{\Gamma(i\kappa)}{\Gamma(\tau)} \text{tr} T_n = \lim_{s \rightarrow \tau} (s - \tau) \sum_{\{P\}, \Gamma_P \neq \{1_2\}} c(P)$$

where the right side is defined as in Theorem 3.13.

Proof. By (2.2), (3.13) and results of [17], we have

$$\lim_{s \rightarrow \tau} (s - \tau) \sum_{j=1, \kappa_j \neq \kappa}^{\infty} h(\kappa_j) \text{tr}_j T_n = 0.$$

By the proof of Theorem 3.13, we have

$$\lim_{s \rightarrow \tau} (s - \tau) \left(d(n) h(-\frac{i}{2}) - c(I) - c(\infty) - \sum_{\{R\}} c(R) \right) = 0.$$

Then the stated identity follows from (2.2), (3.12) and Theorem 3.13. $\square$

Remark. Let h_d be the number of inequivalent classes of primitive indefinite quadratic forms of discriminant d . Siegel [14] proved that

$$(4.1) \quad \lim_{d \rightarrow \infty} \frac{\ln(h_d \ln \epsilon_d)}{\ln d} = \frac{1}{2}.$$

Lemma 4.2. *P is a hyperbolic element of Γ^* with $\Gamma_P \neq \{1_2\}$ if and only if there exists a primitive indefinite quadratic form $[a, b, c]$ of discriminant d such that*

$$P = \frac{1}{\sqrt{n}} \begin{pmatrix} \frac{v - bNu(a, N)^{-1}}{2} & -cNu(a, N)^{-1} \\ aNu(a, N)^{-1} & \frac{v + bNu(a, N)^{-1}}{2} \end{pmatrix}$$

with

$$v^2 - \{dN^2(a, N)^{-2}\}u^2 = 4n.$$

If λ_P is an eigenvalue of P , then

$$\lambda_P - \frac{1}{\lambda_P} = \pm \frac{Nu}{(a, N)} \frac{\sqrt{d}}{\sqrt{n}}.$$

Let

$$P_0 = \begin{pmatrix} \frac{v_0 - bu_0}{2} & -cu_0 \\ au_0 & \frac{v_0 + bu_0}{2} \end{pmatrix},$$

where the pair (v_0, u_0) is the fundamental solution of Pell's equation $v^2 - du^2 = 4$. Then P is $\Gamma_0(N)$ -conjugate to a hyperbolic element $P' \in \Gamma^*$ with $\Gamma_{P'} \neq \{1_2\}$ if and only if P_0 is $\Gamma_0(N)$ -conjugate to P'_0 , where P'_0 is associated with P' similarly as P_0 is associated with P .

Proof. Let

$$P = \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

be a hyperbolic element of Γ^* such that $\Gamma_P \neq \{1_2\}$. Then fixed points r_1, r_2 of P are not rational numbers, and hence Γ_P is the subgroup of elements in $\Gamma_0(N)$ having r_1, r_2 as fixed points. Let $a = C/\mu, b = (D - A)/\mu$ and $c = -B/\mu$, where $\mu = (C, D - A, -B)$. Then $[a, b, c]$ is a primitive quadratic form with r_1, r_2 being the roots of the equation $ar^2 + br + c = 0$. By Sarnak [10], the subgroup of elements in $SL_2(\mathbb{Z})$ having r_1, r_2 as fixed points is generated by the primitive hyperbolic element

$$P_0 = \begin{pmatrix} \frac{v_0 - bu_0}{2} & -cu_0 \\ au_0 & \frac{v_0 + bu_0}{2} \end{pmatrix}$$

where the pair (v_0, u_0) is the fundamental solution of Pell's equation $v^2 - du^2 = 4$.

Since P and P_0 have the same fixed points, we have $A = D - bC/a$ and $B = -cC/a$. Since P belongs to Γ^* and $AD - BC = n$, C satisfies

$$(4.2) \quad \begin{cases} aD^2 - bDC + cC^2 = na \\ a|C, N|C. \end{cases}$$

Let λ_P be an eigenvalue of P . Then it is a solution of the equation $\lambda^2 - \frac{A+D}{\sqrt{n}}\lambda + 1 = 0$. By using $A = D - bC/a$ and $B = -cC/a$, we obtain that

$$(4.3) \quad \lambda_P - \frac{1}{\lambda_P} = \pm \frac{C\sqrt{d}}{a\sqrt{n}}$$

and

$$(4.4) \quad \lambda_P + \frac{1}{\lambda_P} = \frac{1}{\sqrt{n}}(2D - \frac{b}{a}C).$$

Conversely, let a pair (C, D) be a solution of the equation (4.2). Define $A = D - bC/a$ and $B = -cC/a$. Then the matrix

$$P = \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

is a hyperbolic element of Γ^* with $\Gamma_P \neq \{1_2\}$ having the same fixed points as P_0 , and eigenvalues of P satisfies (4.3) and (4.4).

Next, let $v = 2D - b\frac{C}{a}$ and $u_1 = \frac{C}{a}$. Then the equation (4.2) becomes $v^2 - du_1^2 = 4n$ with $N|au_1$. Since $N|au_1$, this equation can be written as

$$(4.5) \quad v^2 - \frac{dN^2}{(a, N)^2} u^2 = 4n.$$

Moreover, we have

$$(4.6) \quad \frac{1}{\sqrt{n}} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \frac{1}{\sqrt{n}} \begin{pmatrix} \frac{v-bNu(a, N)^{-1}}{2} & -cNu(a, N)^{-1} \\ aNu(a, N)^{-1} & \frac{v+bNu(a, N)^{-1}}{2} \end{pmatrix}.$$

Let P'_0 be the primitive hyperbolic element of $SL_2(\mathbb{Z})$ corresponding to $[a', b', c']$. Two forms $[a, b, c]$ and $[a', b', c']$ of the same discriminant are equivalent in $\Gamma_0(N)$ if and only if an element $\gamma \in \Gamma_0(N)$ exists such that $\gamma^{-1}P_0\gamma = P'_0$. $\square$

Lemma 4.3. *Let $[a, b, c]$ be a primitive indefinite quadratic form of discriminant d , and let*

$$P = \frac{1}{\sqrt{n}} \begin{pmatrix} \frac{v-bNu(a, N)^{-1}}{2} & -cNu(a, N)^{-1} \\ aNu(a, N)^{-1} & \frac{v+bNu(a, N)^{-1}}{2} \end{pmatrix}$$

with $v^2 - \{dN^2(a, N)^{-2}\}u^2 = 4n$ be a hyperbolic element of Γ^ with $\Gamma_P \neq \{1_2\}$. Let (v_1, u_1) with $v_1, u_1 > 0$ be the fundamental solution of Pell's equation $v^2 - d_1u^2 = 4$, where $d_1 = dN^2/(a, N)^2$. Then Γ_P is generated by the hyperbolic element*

$$P_1 = \begin{pmatrix} \frac{1}{2} \left(v_1 - b\frac{Nu_1}{(a, N)} \right) & -c\frac{Nu_1}{(a, N)} \\ a\frac{Nu_1}{(a, N)} & \frac{1}{2} \left(v_1 + b\frac{Nu_1}{(a, N)} \right) \end{pmatrix}$$

of $\Gamma_0(N)$.

Proof. Let a, b and c be given as in the proof of Lemma 4.2. Since Γ_P is cyclic, a solution $v_1, u'_1 > 0$ of Pell's equation $v^2 - du^2 = 4$ exists such that Γ_P is generated by

$$P_1 = \begin{pmatrix} \frac{v_1 - bu'_1}{2} & -cu'_1 \\ au'_1 & \frac{v_1 + bu'_1}{2} \end{pmatrix},$$

and hence P_1 is the smallest positive integer power of P_0 among all powers of P_0 belonging to $\Gamma_0(N)$. Since $N|au'_1$, we have $\frac{N}{(a, N)}|u'_1$. Write

$$u'_1 = \frac{Nu_1}{(a, N)}.$$

Then the pair (v_1, u_1) with $v_1, u_1 > 0$ is the fundamental solution of the Pell equation $v^2 - d_1u^2 = 4$, where $d_1 = dN^2(a, N)^{-2}$. The stated result then follows. $\square$

Two quadratic forms $[a, b, c]$ and $[a', b', c']$ are equivalent under $\Gamma_0(N)$ if an element $\gamma \in \Gamma_0(N)$ exists such that

$$\begin{pmatrix} a' & b'/2 \\ b'/2 & c' \end{pmatrix} = \gamma^t \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} \gamma.$$

Lemma 4.4. *Let $[a_j, b_j, c_j]$, $j = 1, 2, \dots, H_d$, be a set of representatives for classes of primitive indefinite quadratic forms of discriminant d , which are not equivalent under $\Gamma_0(N)$. Then we have*

$$\begin{aligned} & \sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) \\ &= 4\sqrt{\pi n} \frac{\Gamma(s - \frac{1}{2})}{N\Gamma(s)} \sum_{d \in \Omega} \sum_{j=1}^{H_d} \sum_u \frac{(a_j, N) \ln \epsilon_{d_1}}{u\sqrt{d}} \left(1 + \frac{d(Nu)^2}{4n(a_j, N)^2}\right)^{\frac{1}{2}-s} \end{aligned}$$

for $\Re s > 1$, where $d_1 = dN^2(a_j, N)^{-2}$ and the sum on u is over all the positive integers u such that $4n + dN^2(a_j, N)^{-2}u^2$ is the square of an integer.

Proof. It follows from (2.1) and Lemma 3.2 that

$$\sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) = 2\sqrt{\pi} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \sum_{\{P\}} \frac{\ln NP_1}{\lambda_P - 1/\lambda_P} \left(1 + \frac{(\lambda_P - 1/\lambda_P)^2}{4}\right)^{\frac{1}{2}-s}$$

for $\Re s > 1$, where $\lambda_P > 1$ is an eigenvalue of P and P_1 is given in Lemma 4.3. Let P be associated with a primitive indefinite quadratic form $[a_j, b_j, c_j]$ as in Lemma 4.2. Then by Lemma 4.2, we have

$$\lambda_P - \frac{1}{\lambda_P} = \frac{Nu}{(a_j, N)} \frac{\sqrt{d}}{\sqrt{n}}.$$

By Lemma 4.3, we have

$$\sqrt{NP_1} = \frac{v_1 + \sqrt{d_1}u_1}{2} = \epsilon_{d_1}.$$

If P' is a hyperbolic of Γ^* with $\Gamma_{P'} \neq \{1_2\}$, and is associated with a primitive indefinite quadratic form $[a_{j'}, b_{j'}, c_{j'}]$ as in Lemma 4.2, then P and P' are $\Gamma_0(N)$ -conjugate if and only if $[a_j, b_j, c_j]$ and $[a_{j'}, b_{j'}, c_{j'}]$ are $\Gamma_0(N)$ -conjugate. Let T be a hyperbolic of Γ^* with $\Gamma_T \neq \{1_2\}$ associated with a primitive indefinite quadratic form $[a, b, c]$ as in Lemma 4.2. If the discriminant of $[a, b, c]$ is not equal to the discriminant of $[a_j, b_j, c_j]$ which is associated with P , then P and T are not $\Gamma_0(N)$ -conjugate. The stated identity then follows. $\square$

The proof for the following lemma is a minor modification of a corresponding lemma in Conrey-Li [2], and is given in Cardon-Li [1].

Lemma 4.5. *Let k be a divisor of N . Then the number of indefinite primitive quadratic forms $[a, b, c]$ with $(a, N) = k$ of discriminant d , which are not equivalent under $\Gamma_0(N)$, is equal to*

$$h_{d_1} \prod_{p^{2l} | (d, k)} p^l \cdot \prod_{p | k} \left(1 + \left(\frac{d}{p}\right)\right)$$

where $d_1 = dN^2/k^2$, where the first product runs over all distinct primes p with p^{2l} being the greatest even p -power factor of (d, k) and the second product is taken over all distinct primes p dividing k .

Theorem 4.6. *Let N be a non-square free positive integer with $(n, N) = 1$. Then we have*

$$\begin{aligned} \sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) &= 4\sqrt{\pi n} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \\ &\times \sum_{k|N} \sum_{m|\frac{N}{k}} \sum_{n|k} \frac{\mu(n)}{n} \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \left(\frac{d}{mn} \right) \frac{h_d \ln \epsilon_d}{u\sqrt{d}} \left(1 + \frac{dk^2 u^2}{4n} \right)^{\frac{1}{2}-s} \end{aligned}$$

for $\Re s > 1$, where the sum on u is over all the positive integers u such that $\sqrt{4n + dk^2 u^2} \in \mathbb{Z}$.

Proof. Let $[a_j, b_j, c_j]$, $j = 1, 2, \dots, H_d$, be a set of representatives for classes of primitive indefinite quadratic forms of discriminant d , which are not equivalent under $\Gamma_0(N)$. By Lemma 4.4, we have

$$\sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) = 4\sqrt{\pi n} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \sum_{d \in \Omega} \sum_{j=1}^{H_d} \sum_u \frac{\ln \epsilon_{d_1}}{u\sqrt{d_1}} \left(1 + \frac{d_1 u^2}{4n} \right)^{\frac{1}{2}-s}$$

for $\Re s > 1$, where $d_1 = dN^2(a_j, N)^{-2}$ and the sum on u is over all the positive integers u such that $\sqrt{4n + d_1 u^2} \in \mathbb{Z}$. Denote $k = N/(a_j, N)$. By Lemma 4.5, we can write the above identity as

$$\begin{aligned} \sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) &= 4\sqrt{\pi n} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \\ (4.6) \quad &\times \sum_{k|N} \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \prod_{p|\frac{N}{k}} \left(1 + \left(\frac{d}{p} \right) \right) \frac{h_{d_1} \ln \epsilon_{d_1}}{u\sqrt{d_1}} \left(1 + \frac{d_1 u^2}{4n} \right)^{\frac{1}{2}-s} \end{aligned}$$

for $\Re s > 1$, where $d_1 = dk^2$. By Dirichlet's class number formula

$$h_{d_1} \ln \epsilon_{d_1} = \sqrt{d_1} L(1, \chi_{d_1})$$

and the identity

$$L(1, \chi_{d_1}) = L(1, \chi_d) \prod_{p|k} \left(1 - \left(\frac{d}{p} \right) p^{-1} \right),$$

we can write (4.6) as

$$\begin{aligned} \sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) &= 4\sqrt{\pi n} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \sum_{k|N} \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \\ &\times \prod_{p|\frac{N}{k}} \left(1 + \left(\frac{d}{p} \right) \right) \prod_{p|k} \left(1 - \left(\frac{d}{p} \right) p^{-1} \right) \frac{h_d \ln \epsilon_d}{u\sqrt{d}} \left(1 + \frac{dk^2 u^2}{4n} \right)^{\frac{1}{2}-s} \end{aligned}$$

for $\Re s > 1$, where the sum on u is over all the positive integers u such that $\sqrt{4n + dk^2u^2} \in \mathbb{Z}$. Since

$$\prod_{p|\frac{N}{k}} \left(1 + \left(\frac{d}{p}\right)\right) \prod_{p|k} \left(1 - \left(\frac{d}{p}\right)p^{-1}\right) = \sum_{m|\frac{N}{k}} \sum_{n|k} \left(\frac{d}{mn}\right) \frac{\mu(n)}{n},$$

we have

$$(4.7) \quad \sum_{\{P\}: P \in \Gamma^*, \Gamma_P \neq \{1_2\}} c(P) = 4\sqrt{\pi n} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \\ \times \sum_{k|N} \sum_{m|\frac{N}{k}} \sum_{n|k} \frac{\mu(n)}{n} \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \left(\frac{d}{mn}\right) \frac{h_d \ln \epsilon_d}{u\sqrt{d}} \left(1 + \frac{dk^2u^2}{4n}\right)^{\frac{1}{2}-s}.$$

Next, we want to show that

$$\sum_{d \in \Omega} \sum_u \left(\frac{d}{mn}\right) \prod_{p^{2l} | (d, \frac{N}{k})} p^l \frac{h_d \ln \epsilon_d}{u\sqrt{d}} \left(1 + \frac{dk^2u^2}{4n}\right)^{\frac{1}{2}-s}$$

is absolutely convergent for $\sigma = \Re s > 1$. Since

$$(4.8) \quad \left| \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \left(\frac{d}{mn}\right) \frac{h_d \ln \epsilon_d}{u\sqrt{d}} \left(1 + \frac{dk^2u^2}{4n}\right)^{\frac{1}{2}-s} \right| \\ \leq \sqrt{N} \sum_{d \in \Omega} \sum_u \frac{h_d \ln \epsilon_d}{u\sqrt{d}} \left(1 + \frac{dk^2u^2}{4n}\right)^{\frac{1}{2}-\sigma}.$$

It is proved in Li [9] that the right side of (4.8) is convergent for $\sigma > 1$, and hence, the right side of the stated identity is absolutely convergent for $\Re s > 1$.

This completes the proof of the theorem. $\square$

Proof of Theorem 1. It is proved at the end of Li [9] that

$$\sum_{d \in \Omega, u} \left| \frac{h_d \ln \epsilon_d}{\sqrt{d}u} \left(1 + \frac{du^2}{4n}\right)^{\frac{1}{2}-\sigma} - (4n)^{\sigma-\frac{1}{2}} \frac{h_d \ln \epsilon_d}{(du^2)^\sigma} \right| \ll \sum_{d \in \Omega, u} \frac{(du^2)^{\frac{1}{2}+\epsilon-1-\sigma}}{u^{1+2\epsilon}} < \infty$$

for $\sigma > 0$. Then it follows from Theorem 4.6 that

$$(4.9) \quad \lim_{s \rightarrow 1/2 + i\kappa} \left(s - \frac{1}{2} - i\kappa\right) \sum_{\{P\}, \Gamma_P \neq \{1_2\}} c(P) = \frac{4\sqrt{n\pi}\Gamma(i\kappa)(4n)^{i\kappa}}{\Gamma(1/2 + i\kappa)} \\ \times \sum_{k|N} \sum_{m|\frac{N}{k}} \sum_{n|k} \frac{\mu(n)}{n} k^{-2i\kappa} \lim_{s \rightarrow 1/2 + i\kappa} \left(s - \frac{1}{2} - i\kappa\right) \sum_{d \in \Omega} \sum_u \prod_{p^{2l} | (d, \frac{N}{k})} p^l \left(\frac{d}{mn}\right) \frac{h_d \ln \epsilon_d}{(du^2)^s}.$$

Theorem 3.13 shows that the function on the right side of (4.9) represents an analytic function in the half-plane $\Re s > 0$ except for a possible pole at $s = 1/2$ and for possible simple poles at $s = 1, \frac{1}{2} \pm i\kappa_j$, $j = 1, 2, \dots$. The stated identity then follows from Lemma 4.1.

This completes the proof of the theorem.

REFERENCES

1. D. A. Cardon and Xian-Jin Li, *A Dirichlet series related to eigenvalues of the Laplacian for congruence subgroups*, Proc. Millennial Conference on Number Theory (Urbana-Champaign, 2000) (to appear).
2. J. B. Conrey and Xian-Jin Li, *On the trace of Hecke operators for Maass forms for congruence subgroups*, Forum Math. **13** (2001), 447–484.
3. J. -M. Deshouillers and H. Iwaniec, *Kloosterman sums and Fourier coefficients of cusp forms*, Invent. Math. **70**, 219–288 (1982).
4. D. A. Hejhal, *The Selberg Trace Formula for $PSL(2, \mathbb{R})$, II, Lecture Notes in Math., no. 1001*, Springer-Verlag, New York, 1983.
5. D. A. Hejhal, *The Selberg trace formula and the Riemann zeta function*, Duke Math. J. **43** (1976), 441–482.
6. H. Iwaniec, *Introduction to the Spectral Theory of Automorphic Forms*, Revista Matemática Iberoamericana, 1995.
7. T. Kubota, *Elementary theory of Eisenstein series*, Halsted Press, New York, 1973.
8. N. V. Kuznecov, *Petersson's conjecture for cusp forms of weight zero and Linnik's conjecture*, Math. USSR Sbornik **39** (1981), 299–342.
9. Xian-Jin Li, *On the trace of Hecke operators for Maass forms*, CRM Proc. and Lecture Notes, Vol. 19, Amer. Math. Soc., Providence, R. I. (1999), 215–229.
10. P. Sarnak, *Class numbers of indefinite binary quadratic forms*, J. Number Theory **15** (1982), 229–247.
11. P. Sarnak, *Arithmetic quantum chaos*, The Schur Lectures (1992, Tel Aviv), Israel Math. Conf. Proc., 8, Bar-Ilan Univ., Ramat Gan (1995), 183–236.
12. A. Selberg, *Harmonic analysis and discontinuous groups on weakly symmetric Riemannian spaces with applications to Dirichlet series*, J. Indian Math. Soc. **20** (1956), 47–87.
13. G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton Univ. Press, 1971.
14. C. L. Siegel, *Über die Classenzahl quadratischer Zahlkörper*, Acta Arith. **1** (1935), 83–86.
15. A. Strömbergsson, *The Selberg trace formula for modular correspondences*, Preprint 1998.
16. E. C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, Second Edition, Oxford, 1948.
17. A. B. Venkov, *Remainder term in the Weyl-Selberg asymptotic formula*, J. Soviet Math. **17** (1981), 2083–2097.
18. G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge Univ. Press, 1952.

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UTAH 84602
E-mail address: xianjin@math.byu.edu